



Factors affecting the availability of data on East African wildlife: the monitoring needs of conservationists are not being met

Inès Moreno^{1,2,3} · Jérôme Marcel Walter Gippet⁴ · Luca Fumagalli¹ · P. J. Stephenson^{1,2}

Received: 28 June 2022 / Revised: 3 October 2022 / Accepted: 20 October 2022 /
Published online: 11 November 2022
© The Author(s) 2022

Abstract

Understanding the status and abundance of species is essential for effective conservation decision-making. However, the availability of species data varies across space, taxonomic groups and data types. A case study was therefore conducted in a high biodiversity region—East Africa—to evaluate data biases, the factors influencing data availability, and the consequences for conservation. In each of the eleven target countries, priority animal species were identified as threatened species that are protected by national governments, international conventions or conservation NGOs. We assessed data gaps and biases in the IUCN Red List of Threatened Species, the Global Biodiversity Information Facility and the Living Planet Index. A survey of practitioners and decision makers was conducted to confirm and assess consequences of these biases on biodiversity conservation efforts. Our results showed data on species occurrence and population trends were available for a significantly higher proportion of vertebrates than invertebrates. We observed a geographical bias, with higher tourism income countries having more priority species and more species with data than lower tourism income countries. Conservationists surveyed felt that, of the 40 types of data investigated, those data that are most important to conservation projects are the most difficult to access. The main challenges to data accessibility are excessive expense, technological challenges, and a lack of resources to process and analyse data. With this information, practitioners and decision makers can prioritise how and where to fill gaps to improve data availability and use, and ensure biodiversity monitoring is improved and conservation impacts enhanced.

Keywords Biodiversity monitoring · Data availability · East Africa · Global biodiversity information facility · IUCN red list of threatened species · Living Planet Index

Communicated by Alison Nazareno.

✉ P. J. Stephenson
StephensonPJ@gmail.com

Extended author information available on the last page of the article

Introduction

An understanding of the status and abundance of species, their habitats, the threats and pressures they face, and the progress of work undertaken for their conservation is essential for effective project management and decision-making (Robinson et al. 2018; Hu et al. 2019; Stephenson and Stengel 2020). However, for a variety of reasons, species monitoring is largely inadequate and often neglected, meaning the necessary data are unavailable to decision makers (Amano et al. 2016; Stephenson et al. 2017a, 2022). Moreover, scientific knowledge of species is constrained by the taxonomic and spatial biases of the available data (Amano et al. 2016; Troudet et al. 2017; dos Santos et al. 2020).

Taxonomic biases in biodiversity research and monitoring at the global level tend to result in an under-representation of invertebrates in data sets (Leather 2009; Hochkirch et al. 2021) and an over-representation of vertebrates, especially mammals and birds (Bonnet et al. 2002; McRae et al. 2017; Moussy et al. 2022). Geographic biases mean the tropical regions housing the most biodiversity and the most threatened species are the least studied (Pimm et al. 2006; Amano and Sutherland 2013; Titley et al. 2017). Reasons for this inequality are diverse and poorly understood (Stephenson et al. 2017a, 2022; Stephenson 2019; Hochkirch et al. 2021) but may be a key factor in the low use of data in national biodiversity reports (Bubb et al. 2011).

It is necessary to understand and address the gaps and biases in biodiversity knowledge if we are to enhance conservation management. The urgency is particularly acute in Africa where a large proportion of the population is directly dependent on ecosystem services for their livelihoods, yet increasing habitat loss from the expansion of agricultural land and urban areas is exacerbating biodiversity loss (Craigie et al. 2010; Seto et al. 2012; IPBES 2018).

The aim of our research was therefore to conduct a case study in East Africa to answer the question: What are the taxonomic and geographic gaps and biases in the biodiversity data available for species considered priorities and how are they affecting conservation action? The project set out to identify trends in data availability and place them in the context of recognised conservation priorities at national, regional and global levels. We conducted an analysis of global databases to identify overall trends in data availability for the region, and to discover to what extent regional biases reflected global biases. However, in order to gain an understanding of data user needs and the root causes and consequences of data gaps, we conducted a survey of practitioners and policymakers. Our ultimate aim was to understand the factors blocking the flow of biodiversity data to conservation decision makers in East Africa.

We focused on assessing the biodiversity data needs and challenges in eleven countries in East Africa: Burundi, Comoros Islands, Ethiopia, Kenya, Madagascar, Malawi, Mozambique, Rwanda, Somalia, Tanzania and Uganda. This region was selected as it has a wide range of habitats (including rainforests, mangroves, mountains, freshwater lakes, coral reefs and seagrass beds), and several priority ecoregions (e.g. African Rift Lakes, Coastal East Africa, Coastal East Africa Marine, and Madagascar; Olson and Dinerstein 2002) and associated biodiversity hotspots (Eastern Afrotropical, Coastal Forests of Eastern Africa, and Madagascar and the Indian Ocean; Mittermeier et al. 2004). As a result, the region has been the focus of conservation efforts by many national governments and national and international conservation agencies dating back several decades (Huxley 1961). Although there are increasing efforts to identify global

floral and fungal conservation priorities (e.g. Bachman et al. 2018; Gonçalves et al. 2021), governments and NGOs in East Africa primarily focus their attention on their faunas and so our study looked at animal species only.

Many biodiversity monitoring systems fail to take account of user needs (Stephenson et al. 2015b, 2017b). While exploring data on threatened species can be useful, the level of extinction risk is not the same as the level of priority for conservation action (Fitzpatrick et al. 2007; Mace et al. 2007; Le Berre et al. 2019). The most threatened species are often the most difficult and the most expensive to conserve (Le Berre et al. 2019) and some taxa will be priorities for conservation action even if they are not listed as threatened (IUCN 2012). Therefore, in this study regional data needs were determined by defining priority species from national laws and governmental, inter-governmental and NGO conservation plans. Data availability and access challenges were identified by assessing East African data in three global databases commonly used to monitor delivery of national contributions to global biodiversity goals: the International Union for Conservation of Nature (IUCN) Red List of Threatened Species (IUCN 2021), the WWF/ZSL Living Planet Index (Loh et al. 2005; Living Planet Index 2021) and GBIF—the Global Biodiversity Information Facility (GBIF 2021). We assessed the results against socio-economic factors and surveyed data users to better understand the reasons and consequences of data access issues in the region.

Methods

Data collection

Threatened species

Threatened animal species in the 11 countries were identified through the IUCN Red List of Threatened Species (hereafter the IUCN Red List; Fig. 1). The IUCN Red List (IUCN 2021) is an inventory of the global conservation status of biodiversity (animals, plants and fungi). Using a set of quantitative criteria, it assesses the risk of extinction of species, ranging from Least Concern (for species that are still abundant in the wild and under little threat) to Extinct (IUCN 2012). Threatened species are those assessed to be in categories Vulnerable, Endangered and Critically Endangered. Data on range, population size, habitat and ecology, use and trade, threats, and conservation actions are also listed in the database. Data downloaded for this study included: scientific name, taxonomy, Red List category, population trend (stable, decreasing, increasing or unknown), assessment ID, assessment date, year published, system (e.g. marine, terrestrial, freshwater), countries where each species was present, and potential threats. (References and DOIs for the Red List data downloaded can be accessed in the supplementary material Appendix A, Tables S1 and S2 respectively). Data files containing each species' countries and threats were downloaded separately and modified using the Python programming language (version 3.6; VanRossum and Drake 2009) to allow for better visualisation of the data and to make it easier to conduct analyses. The total numbers of species listed in the IUCN Red List by target country and species class in all 11 target countries were also recorded for comparison with the numbers of threatened and priority species (see below) obtained for each target country and species class in the study area.




Fig. 1 Map of Africa showing the relative number of threatened animal species in each East African country studied: Burundi, Comoros Islands, Ethiopia, Kenya, Madagascar, Malawi, Mozambique, Rwanda, Somalia, Tanzania and Uganda. The darker the colour, the greater the number of threatened species

Priority species

While many people consider threatened species to be important, data are more likely to be collected for those that are considered a priority by key stakeholder groups. Therefore, we defined a priority species as a threatened species identified for protection by law or by action by a government, international convention or non-governmental organisation (NGO) (Fig. 2). We focused on priority species because most conservationists do not need to monitor all species in every taxon; it is infeasible (especially for very speciose groups like invertebrates) and unnecessary, since data are only required for species that are of concern or are the focus of conservation action (i.e., priority species).

After identifying threatened species, we identified which ones were priorities through a review of national laws and strategies, Multilateral Environmental Agreements (MEAs), reports to the Convention on Biological Diversity (CBD; www.cbd.int), and NGO strategies. Each country's National Biodiversity Strategy and Action Plan (NBSAP) was reviewed, as were National Reports to the CBD. Lists of species protected by law were found on ECOLEX (IUCN, UNEP and FAO 2021) and the FAOLEX Database (FAO




Fig. 2 Workflow used in the present study. The number of threatened animal species was first identified from the IUCN Red List. Of these, species protected by the governments (GOs), species listed on International Conventions (CI), i.e. on the Convention on International Trade in Endangered Species of Wild Fauna and Flora (CITES) and on the Convention on the Conservation of Migratory Species of Wild Animals (CMS), as well as species prioritised by NGOs operating in the selected countries were listed as "priority" species. Data of these priority species was then collected from the Global Biodiversity Information Facility (GBIF) and Living Planet Index (LPI). The number of priority species for which data in each of the two databases is shown in the yellow rectangles. In parallel, a survey of practitioners and policy makers was conducted

2021). Both protected and partially protected species were considered priorities. Names and dates of the laws found for each country can be found in supplementary material (Appendix A, Table S3). The protected species lists of two international conventions, the Convention on International Trade in Endangered Species of Wild Fauna and Flora (CITES) and the Convention on the Conservation of Migratory Species of Wild Animals (CMS), were also used to identify priority species in target countries (Fig. 2). All 11 countries studied are parties to both conventions and so, by definition see their goals and priorities as national priorities. The lists of species protected by the conventions were downloaded from the Species+ website (UNEP-WCMC 2021) for CITES appendices I, II and III and CMS appendices I and II.

Through web searches we identified the main international NGOs operating in the target region and for which at least some information was available online, which included: African Wildlife Foundation (AWF), Alliance for Zero Extinction (AZE), Conservation International,

Re:wild (formerly Global Wildlife Conservation), Wildlife Conservation Society (WCS), World Wide Fund for Nature (WWF), and the Zoological Society of London (ZSL). Priority species per country for these NGOs were then identified from their respective websites in April and May 2021. No priority species were found for Conservation International. A list of priority species per country was then created. For a species to be considered a priority it had to be: i) present in the country, ii) considered a priority by the government of the country or by one of the NGOs working in that country or on the country's list of one of the two international conventions (CITES or CMS).

Data availability

We then assessed data availability for priority species by interrogating the Global Biodiversity Information Facility (GBIF; www.gbif.org, and the Living Planet Index (LPI; www.livingplanetindex.org) (Fig. 2). (Citations for each data download can be found in the supplementary material Appendix A, Table S4). These global databases bring together several types of biodiversity data and were chosen for this study as they are the most extensive biodiversity databases currently available, and are used to measure indicators by the Convention on Biological Diversity (CBD; Brooks et al. 2015; McRae et al. 2017), of which all of the East African countries are signatories. To assess whether there had been increased monitoring in response to the CBD adoption of the Strategic Plan for Biodiversity 2011–2020 (Convention on Biological Diversity 2010), the number of data collected since 2010 were analysed separately for GBIF and LPI databases.

GBIF is a global database of species presence records collected from multiple sources ranging from museum specimens to geo-referenced photographs and records from amateur naturalists. It is linked to, and holds data from, other global databases, including eBird (<https://ebird.org>). In December 2021, 1,902,873,733 occurrences were recorded in the database. GBIF records of East African priority species were downloaded via the GBIF website (GBIF 2021) in August and October 2021 (Fig. 2). As differences in the species number of occurrences were observed when the 'country' filter was applied directly on the GBIF database website or after downloading the species occurrences, all occurrences found for every species were downloaded and subsequently sorted by country. Data extracted included the number of occurrences recorded by country and the date of the last record and the basis of the last record (e.g. "human observation", "preserved specimen"). The final GBIF data file can be accessed in supplementary Table S4, Appendix A.

The LPI is a database of population trends of vertebrate species managed by the Zoological Society of London, which enables the assessment of changes in biodiversity over time. The database holds time-series data for over 27,000 populations of more than 4300 species. These population trend data are collated from different sources such as published literature, grey literature, government reports and online databases. Data on population trends of priority species were downloaded from the LPI data portal (Fig. 2; Living Planet Index 2021). Data extracted included the number of time series recorded by species by country and the date of the last record. The final LPI data file can be accessed in supplementary Table S5, Appendix A.

Socio-economic factors: international tourism receipts, gross domestic product and political stability

Some studies (e.g., Moussy et al. 2022) have shown that socio-economic variables may influence biodiversity data availability, and so economic data on target countries were collected. A web search investigated potential socio-economic data that might be of relevance in understanding biodiversity trends in East Africa. The most complete data sets found were for three socio-economic variables: gross domestic product (GDP), international tourism receipts and political stability.

GDP figures in billions of US dollars were obtained from the International Monetary Fund database (International Monetary Fund 2021). Data on international tourism receipts for each target country from 2014 to 2018 were obtained from the World Tourism Organization (UNWTO) Tourism Highlights Reports from 2016 to 2020 (UNWTO 2021). Somalia had no data on international tourism receipts. As these values fluctuate from year to year, an average of GDP (in millions of US dollars) and international tourism receipts (in millions of US dollars) from 2014 to 2018 was calculated for each country. Country population size data were collected from UN data (www.data.un.org) to take into account the tourism income and the GDP per capita in the analyses. The Political Stability Index, measured as “perceptions of the likelihood of political instability and/or politically-motivated violence, including terrorism”, of each country from 2014 to 2018 were obtained from the World Bank database (www.databank.worldbank.org). These country-specific socio-economic data can be found in supplementary material Appendix A Table S6.

Survey of data users

An online survey of conservation practitioners and policy makers was conducted to understand data user needs in East Africa and the root causes and consequences of data gaps. Questionnaire surveys have been used extensively in conservation biology (e.g., Tenopir et al. 2011; Chen et al. 2019; Danovaro et al. 2020) and allow direct input from stakeholders (Sanders et al. 2021) to help provide answers that may not be found in the literature, where data is scarce or missing (Meyer and Booker 2001).

While the global databases interrogated primarily provide data of use in measuring species presence (GBIF) or population trends (LPI), conservation managers ultimately need data to measure a much broader range of indicators on the state of biodiversity, the pressures species and habitats face and the conservation responses made to reduce pressures (Tittensor et al. 2014; Stephenson 2019; Stephenson et al. 2022). Therefore, in addition to surveying user needs for data of the type captured in the target global databases, we also included questions around other indicators to gain a more complete understanding of the issues. Forty data types were identified for inclusion in the survey, based on variables used for Red List assessments (IUCN 2012) and common metrics for project management (Table 1).

A confidential online survey for practitioners and policy makers was designed using Qualtrics survey software (Qualtrics, Provo, UT) and emailed to 65 conservation agencies: the environmental authority of each target country and 54 NGOs working in at least one of the countries. The survey was open from 8 September to 11 October 2021. Twenty-one questions were asked (in English and French) to obtain information on respondents' conservation projects (recent, current or upcoming), the data they need to carry out their

Table 1 The 40 types of data assessed in the survey

Data types

Population parameters

1. Presence/absence
2. Number of individuals (density/abundance)
3. Species range/distribution
4. Number of populations

Demographic parameters

5. Reproductive rates
6. Mortality rates
7. Genetic parameters
8. Movement patterns
9. Health/physiological condition

Habitat

10. Habitat extent/cover
11. Habitat quality

Threats

12. Poaching levels
13. Levels of trade in the species or its body parts
14. Pollution affecting the species
15. Invasive species impacting the species
16. Levels of human-wildlife conflict involving this species
17. Development in species range
18. Transportation or service corridors in species range
19. Agriculture or aquaculture in species range
20. Energy production and mining in species range
21. Natural systems modification in species range (e.g. dams, fires)
22. Climate change and severe weather events

Associated species

23. Prey species
24. Predator species
25. Competitor species
26. Parasites/pathogens

Already ongoing conservation efforts

27. Site/area protection
28. Site/area management
29. Habitat and natural resources protection
30. Habitat and natural processes restoration
31. Trade controls
32. Species re-introductions
33. Ex-situ conservation (e.g. captive breeding, genome resource bank, ...)
34. Strategies to mitigate human-wildlife conflicts
35. Invasive alien species management
36. Education and awareness initiatives

Table 1 (continued)

Data types

-
- 37. Law and policy development
 - 38. Law and policy enforcement (including anti-poaching)
 - 39. Livelihood, economic and other incentives
- Socio-economic and cultural value of the species
- 40. Socio-economic and cultural value of the species
-

projects, data collection, difficulties encountered in accessing data, the impact of a lack of data on their projects, and data sharing.. The questionnaire survey is available in supplementary Table S7, Appendix A.

Data analyses

All data analyses were carried out using the R statistical programming language version 4.1.1 (R Core Team 2021). First, we assessed the taxonomic bias in data availability. Using Pearson’s Chi-Squared tests (“prop.test” function in *stats* package), we compared the proportions between invertebrates and vertebrates, and the proportions among vertebrates for (i) the proportion of threatened species considered as a priority, (ii) the proportion of priority species with IUCN population trends data, (iii) the proportion of priority species with GBIF data and (iv) the proportion of priority species with GBIF data since 2010 among vertebrate classes. We did not test for differences in proportion of priority species with LPI data because this data was lacking for most taxonomic groups (except birds and mammals). We applied Bonferroni corrections to account for multiple comparisons.

Secondly, we tested if the number of species considered as a conservation priority was linked to tourism receipts per capita, GDP per capita or political stability. Five different models were performed using the *glmmTMB* R package (v1.1.2.3; Brooks et al. 2017), each using a different measure of the number of priority species by country as response variable: (i) the number of priority species in each country, (ii) the number of priority species per country having population trend data in the IUCN Red List, (iii) the number of priority species per country for which data are available in GBIF, (iv) in GBIF since 2010, (v) in LPI, or (vi) in LPI since 2010. Because these response variables are counts data, we used generalized linear models (GLMs) with Poisson or negative binomial distribution (negative binomial distribution was used if there was overdispersion in the model’s residuals). The validity of the models was assessed using the R package *DHARMA* (v0.4.4; Hartig 2021). The final models can be found in Appendix A, Table S8. Explanatory variables (tourism receipts per capita, GDP per capita and political stability indexes) were log-scaled in the models. The effect of each explanatory variable was tested using an analysis of variance (“Anova” function in *car* package, v3.0–11; Fox and Weisberg 2019).

Thirdly, responses to the survey of practitioners and policy makers were analysed. The relationship between the importance of data for successful conservation projects (i.e., the proportion of respondents who think the data are important) and the difficulty of accessing the same data (i.e., the proportion of respondents who think the data are difficult to access) was tested. As our response variables are percentages, binomial models were computed using the *glmmTMB* R package (v1.1.2.3; Brooks et al. 2017). The 40 data types (Table 1) were divided into seven categories: population parameters, demographic parameters,

habitat, threats, associated species, already ongoing efforts, and socio-economic and cultural value of the species. These categories were accounted for in the analyses as a random effect. Post hoc multiple comparisons (with Bonferroni correction) were conducted to assess the differences between groups (“*cld*” function, package *multcomp*, version 1.4–17, Piepho 2004). The validity of our models was assessed using R package *DHARMA* (v0.4.4; Hartig 2021) and the effect of the explanatory variable was tested using an analysis of variance (“*Anova*” function, *car* package, v3.0–11; Fox and Weisberg 2019).

Results

Data availability for threatened and priority species

In the 11 study countries, data on 11,071 animal species were available in the IUCN Red List, of which 1674 species (15.1%) were assessed as threatened. There are at least 55,683 unique animal species names recorded in the region (GBIF.org [28 August 2022] GBIF Occurrence Download), meaning that the 11,071 species assessed in the Red List represents about 19.9% of the total fauna. Of the 1674 threatened species, 681 were considered priority species, with 293 species protected by governments, 364 species listed as protected on international conventions (CMS and CITES), and 336 considered a priority by one or more NGOs (Fig. 2, 3 and 4).

Taxonomic bias in data availability

Four of the nine invertebrate classes, Anthozoans (sea anemones and corals), Holothurians (sea cucumbers), insects and Malacostracan crustaceans (crabs, lobsters, shrimps, etc.), had priority species, but this represented only 79 (11.6%) of the total priority species for the region. In comparison, all seven vertebrate classes with threatened species in East Africa included at least one priority species (Fig. 3). The proportion of threatened species considered as a priority was significantly higher for vertebrates (0.476) than for invertebrates (0.193) (Pearson’s Chi-Squared test, $\chi^2_1 = 101.22$, $p < 0.00001$). However, Holothurians (sea cucumbers) had the highest proportion (90%) of priority species among threatened species than any vertebrate or invertebrate class. The proportion of threatened species considered as priority varied among vertebrate classes (Pearson’s Chi-Squared test, $\chi^2_6 = 417.03$, $p < 0.0001$). Most threatened Chondrichthyes species (cartilaginous fishes) were listed as priorities while Actinopterygians species (ray-finned fishes) were rarely listed as priorities (see Fig. 4 and supplementary Table S9 for details).

The IUCN Red List provides an assessment of the population trends for some species. The proportion of threatened species having population trends data on the IUCN Red List was significantly higher for vertebrates (0.740) than for invertebrates (0.413) (Pearson’s Chi-Squared test, $\chi^2_1 = 145.59$, $p < 0.0001$). The same applies to priority species with a proportion of vertebrates and invertebrates of 0.875 and 0.430 respectively (Pearson’s Chi-Squared test, $\chi^2_1 = 92.24$, $p < 0.0001$). The proportion of priority species having population trends data on the IUCN Red List varied among vertebrate classes (Pearson’s Chi-Squared test, $\chi^2_6 = 39.94$, $p < 0.0001$). Birds had a higher proportion of priority species and reptiles and Actinopterygians (ray-finned fishes) had a lower proportion of priority species with population trends data on the IUCN Red List than most other classes of vertebrates (see Fig. 4 and supplementary Table S9 for details).




Fig. 3 Number of threatened and priority animal species per country. The numbers in brackets are the total number of animal species per country assessed by the IUCN Red List (critically endangered, endangered, vulnerable, lower risk; conservation dependent, near threatened, last concern, data deficient). Left panel: Number of threatened species (in light green) according to the IUCN Red List and of these the number of species considered as priorities (in dark blue) according to international conventions, governments and NGOs. Right panel: number of priority species (dark blue) per country, number of priority species with population trend estimates in the IUCN Red List (red), and the number of priority species that have (i) ≥ 1 occurrence in GBIF (green), (ii) ≥ 1 occurrence in GBIF since 2010 (hatched green), (iii) population trends data in LPI (yellow) and (iv) in LPI since 2010 (hatched yellow) per country. x axis of the right panel is log scaled

Of the 681 priority species, 609 (89.4%) had one or more occurrence records in GBIF, 561 (82.4%) had population trend data on the IUCN Red List and 49 (7.2%) had population trend data in the LPI. When Data were considered from 2010 onwards (during implementation of the CBD Strategic Plan for Biodiversity 2011–2020), occurrence data were available for 397 priority species in GBIF (a 35% reduction) and 11 priority species in the LPI (a 78% reduction).

Data existed in GBIF for 554 out of 602 (92.0%) priority vertebrate species and 55 out of 79 (69.6%) priority invertebrate species. However, there are data for only 379 (63%) of priority vertebrates and 18 (22.8%) of priority invertebrates since 2010. The proportion of priority species with data in GBIF was significantly higher for vertebrates (0.920) than for invertebrates (0.696) (Pearson’s Chi-Squared test, $\chi^2_1 = 34.75, p < 0.0001$). The same applies to priority species with data in GBIF since 2010 (Pearson’s Chi-Squared test, $\chi^2_1 = 44.72, p < 0.0001$). Among vertebrate classes, there was no significant differences between the proportion of priority species with data in GBIF. However, when focusing on data produced since 2010, the proportion of priority species with data on GBIF varied among vertebrate classes (Pearson’s Chi-Squared test, $\chi^2_6 = 78.87, p < 0.0001$). Birds had




Fig. 4 Number of threatened and priority animal species per class in East Africa. The numbers in brackets are the total number of extant animal species per class in the 11 target countries assessed by the IUCN Red List (critically endangered, endangered, vulnerable, lower risk; conservation dependent, near threatened, last concern, data deficient). Left panel: Number of threatened species (in light green) per class in the 11 target countries according to the IUCN Red List and of these the number of species considered as priorities (in dark blue) according to international conventions, governments and NGOs. Right: number of priority species (dark blue) per class in the 11 target countries, the number of priority species with population trend estimates in the IUCN Red List (red) and the number of priority species that have (i) ≥ 1 occurrence in GBIF (green), (ii) ≥ 1 occurrence in GBIF since 2010 (hatched green), (iii) population trends data in LPI (yellow) and (iv) in LPI since 2010 (hatched yellow) per class in the 11 target countries. x axis of the right panel is log scaled

more priority species than any other vertebrate classes while amphibians were the less represented vertebrate class on GBIF since 2010 (see Fig. 4 and supplementary Table S9 for details).

Geographic and socio-economic bias in data availability

The number of priority species per country was positively correlated with per capita international tourism receipts (Table 2; Fig. S1), as was the number of priority species having population estimates in the LPI and having ≥ 1 occurrence records in GBIF and in GBIF since 2010. The number of priority species with LPI data was also positively correlated

Table 2 Assessment of data against socio-economic factors: tourism receipts, GDP and political stability

Correlation	Tourism receipts		GDP		Political stability	
	Chisq (Df = 1)	<i>p</i> value	Chisq (Df = 1)	<i>p</i> value	Chisq (Df = 1)	<i>p</i> value
Number of priority species	5.174	0.023	0.761	0.383	0.163	0.686
Nb of priority species having population trends in the IUCN RL	0.961	0.327	0.423	0.516	0.148	0.700
Nb priority species having ≥1 occurrence in GBIF	6.676	0.010	1.834	0.176	3.536	0.060
Nb priority species having ≥1 occurrence in GBIF since 2010	5.348	0.021	1.263	0.261	1.436	0.231
Nb priority species having population trend data in LPI	11.232	0.001	4.315	0.038	1.839	0.175
Nb priority species having population trend data in LPI since 2010	3.459	0.063	0.956	0.328	2.058	0.152

Statistically significant values (*p* < 0.05) are marked in bold

with per capita GDP. There was no correlation between each country’s Political Stability Index and the number of priority species in each country or database (Table 2).

Data users’ perceptions

We received 33 completed responses to the online survey of species conservation experts (a response rate of 51%). These included three responses from environmental authorities (in the governments of Comoros, Malawi and Rwanda), and 30 responses




Fig. 5 Factors affecting data accessibility and usability

from NGOs (see acknowledgements). Based on responses received, the top five factors affecting the accessibility and usability of data were excessive expense, technological challenges, lack of capacity, the incompleteness of data, and the poor quality of available data (Fig. 5). The top five types of data required by decision makers related to species abundance, habitat extent, habitat quality, levels of human-wildlife conflict and conservation responses for site protection (Fig. 6a). Difficulty in accessing data was positively correlated with the importance of the data for conservation projects (Fig. 6a) and differed by data type (Fig. 6b) (binomial GLM: $\chi^2_1 = 4.8, p = 0.028$ and $\chi^2_6 = 27.41, p = 0.0001$ respectively). The data most important for respondents was the most difficult for them to access.




Fig. 6 Relationship between the difficulty of access to data and the importance of these data for carrying out conservation projects. **a** Correlation between the importance of data and the difficulty to access these data. **b** Differences between data group means ranked with letters. Group means sharing a letter are not significantly different. Data type group 1: population parameters, group 2: demographic parameters, group 3: habitat, group 4: threats, group 5: associated species, 6: already ongoing conservation efforts, group 7: socio-economic and cultural value of the species

Discussion

Priority species

This study is unique in that it considered not just the availability of biodiversity data but the availability of data for species that have been defined as priorities by governments and the NGOs supporting them. “Allocating resources solely to the most endangered species will typically not minimize the number of extinctions in the long-term, as this does not account for the risk of less endangered species going extinct in the future” (Wilson et al. 2011). Therefore, we looked at not just the most threatened animals in East Africa, but those considered priorities by the conservation community. However, identifying national priorities was often a difficult task. In general, countries did not specify priority species in their NBSAPs or in their reports to the CBD. NGO priorities were also sometimes difficult to identify from the organisations’ websites, with details buried in hard-to-find webpages or inaccessible plans and reports. The conservation organisations often gave only examples of priorities instead of full lists. Other studies in Africa have found that conservation plans need taxonomic and spatially explicit details to facilitate effective delivery and monitoring (e.g., Balmford 2003; Stephenson and Ntiamoa-Baidu 2010). In the future, targeted conservation action and monitoring might be improved if governments and NGOs specified, better communicated and reported on the taxa they focus their conservation attention on.

Taxonomic data gaps and biases

As expected, across East Africa more vertebrate species are considered priorities than invertebrate species. However, Holothurians (sea cucumbers) and Anthozoans (corals and their relatives) had relatively high proportions of threatened species considered priorities. This may reflect the value of these taxonomic groups to national and local economies due to the exploitation of sea cucumbers for food and of coral reefs for fisheries and tourism. East Africa is also important for many threatened and priority marine vertebrate species such as sea turtles, dugongs, dolphins, rays and many fish species (Sievers et al. 2019), and National Reports to the CBD of the Comoros, Ethiopia, Kenya, Madagascar and Tanzania (Convention on Biological Diversity 2021) state that actions are being taken to meet the Aichi Target 10 on ocean ecosystems.

Although data were available in GBIF for more than two thirds of the regional priority invertebrates, this was significantly less across taxa than for vertebrates. The findings in East Africa reflect historic global trends in taxonomic data biases towards vertebrates (Loh et al. 2005; Troudet et al. 2017), partly reflecting an over-representation of mammals and birds within vertebrates research and monitoring (Bonnet et al. 2002; McRae et al. 2017; Christie et al. 2021; Moussy et al. 2022). However, even within these classes, there are gaps with, for example, less data on African small mammals than African large mammals (Stephenson et al. 2021).

Actinopterygians (ray-finned fishes) were well represented in GBIF and the LPI which may be explained partly by the fact that East African countries are dependent on fisheries for food and economic income (Garibaldi 2014), and exploited fish stocks are likely to be better monitored (McRae et al. 2017). The trend for significantly fewer data on invertebrates, especially insects, is seen by many as worrying given the importance of these species-rich taxa

for a variety of ecosystems and ecosystem services (Gascon et al. 2015; Cardoso and Leather 2019).

There were fewer data on priority vertebrate species in the LPI than in GBIF. This may partly reflect the fact it is easier to collect an occurrence record for a species (e.g., using museum collection records or citizen science observations) than it is to conduct a survey of the population with enough detail to calculate trends in abundance over time. Nonetheless, time-series data were found for only 11 East African priority species (mostly mammals) since 2010. Previous assessments of the LPI have acknowledged its geographic biases, with the majority of data collected in temperate regions rather than high biodiversity tropical regions (Collen et al. 2009). Reasons for low data input may include the lack of research conducted to obtain data on population trends and a lack of data sharing when data are collected (McRae et al. 2017). Eight people surveyed in this study explicitly stated that they were not aware of the existence of the LPI database, reflecting other findings that some practitioners are not aware of global databases or confident of the relevance of database content (Bowles-Newark et al. 2015). This further confirms the need to raise awareness of global databases at national level.

Troudet et al. (2017) suggested that public opinion guides biodiversity data gathering. Since many of the European and North American cultures that provide a high proportion of the resources for global databases usually perceive vertebrates such as large mammals to be the most charismatic species (Colléony et al. 2017; Krause and Robinson 2017; Albert et al. 2018); it may not be surprising that vertebrates dominate those databases. There is also often an assumption that, if larger, wider-ranging species such as vertebrates are conserved, then smaller species in the same habitats, such as invertebrates, will also be conserved. This may explain some of the data collection biases. However, the use of such surrogates has its pitfalls (Cardoso et al., 2011) and may mean that taxa not monitored decline without us knowing. This is why calls have been made to enhance data collection for neglected taxa (Hochkirch et al., 2021; Stephenson et al., 2022).

Of course, the presence of data in a global database—whether occurrence records or trends in populations—does not in itself mean the data are useful for monitoring or reporting. The records need to be appropriate for the users (e.g., accessible, timely, relevant, easily understood). However, several studies have shown the potential value of disaggregating global data sets to obtain national trends (Han et al. 2014; Stephenson et al. 2015a). “Global datasets cannot always replace local or national data” (Bowles-Newark et al. 2015) but certain indicators such as population trends lend themselves to national analyses (Stephenson et al. 2015a) and the LPI has been used to create several national reports (e.g., van Strien et al. 2016; WWF-Canada 2020). In future, conservationists in East Africa need to explore further the use of global data sets to enhance their monitoring and reporting on biodiversity. If they are to ramp up the monitoring of their contributions to the CBD’s post-2020 global biodiversity framework, countries will not only need more and better data on species occurrence and population trends, but also on the pressures faced by species and the conservation responses made (Tittensor et al. 2014; Stephenson 2019), including the links between ecological systems and social and governance systems (Mastrángelo et al. 2019). However, our findings allow us a proxy measure of the availability of useful data at a global level and it suggests there remain some significant gaps in East Africa.

Filling data gaps

The absence of data on some priority species may reflect a lack of data collection or lack of sharing collected data. It may also reflect the fact only one in five countries worldwide uses

national indicators based on those recommended by CBD (Bhatt et al. 2020). Conservation project managers often report more on activities and outputs rather than impacts (Stephenson 2019), and one review found only 19% of species-focused conservation projects submitted data on population trends (Badalotti et al. 2022). This suggests a more fundamental need to improve impact monitoring in conservation projects in East Africa and beyond.

More data need to be collected on many species considered as conservation priorities, especially invertebrates, amphibians and reptiles, as well as marine species. Efforts are already underway to incorporate data on invertebrates into the LPI, as has been started with European grassland butterflies (WWF 2020). But these efforts will only succeed if more fundamental improvements are made in species monitoring at national level.

As proposed by Hochkirch et al. (2021), measures and tools to enter research results systematically into global databases should be put in place, such as the obligation by scientific journals to share population trend or occurrence data through databases such as GBIF and the LPI. To motivate such behaviour, data contributed to global databases needs to be recognised as a credible publication (Costello et al. 2013) with mechanisms put in place for data citation to be comparable to other scientific publications (Costello 2009). The IUCN Red List already allows species assessments to be saved and allocated a digital objective identifier which therefore promotes the sharing of data by making it a citable publication. Other databases should follow suit.

Citizen science, the engagement of people without scientific training in the collection of data, has traditionally been focused on Western Europe and North America, but in recent years has expanded in Africa (Wotton et al. 2020; Stephenson et al. 2021). While this may provide additional opportunities for monitoring (see Chandler et al. 2017), efforts will need to be made to ensure data-deficient species are included (Theobald et al. 2015; Troudet et al. 2017). As well as exploring options for citizen science, more effort is needed to tap into indigenous knowledge (Sitati and Ipara 2012); many local communities in East Africa will know better than universities, NGOs or governments if a certain species occurs near them or not.

Addressing factors affecting data availability and use

The importance of biodiversity data for effective biodiversity conservation management has been well documented (Reichman et al. 2011; Stephenson and Stengel 2020) but our study shows for the first time that the data considered by practitioners as the most useful are the data that are most difficult to access. The data needed most are related to species abundance and the extent and quality of habitat (Fig. 6), which are common measures of biodiversity state. The three main challenges identified for data access and use revolved around inadequate resources, tools and capacity, issues raised in other studies (Amano and Sutherland 2013; Thapa et al. 2014; Tittensor et al. 2014; Stephenson et al. 2017a, 2021; Rounsevell et al. 2020).

We used three global databases (IUCN Red List, GBIF and LPI) as proxies for the taxonomic and geographic biases of biodiversity data relevant to East Africa. However, we note that other databases (at global and local levels) are needed to support countries with tracking other indicators not related to species presence or population trends. Therefore, a follow up study would be useful to determine the availability and biases of other data rated as important, such as habitat cover, habitat quality and human-wildlife conflict.

It is notable that, while the databases reviewed are openly and freely accessible, two of the greatest challenges to data use were excessive expense and a lack of resources to

process and analyse data. This suggests that open-access databases have associated costs for users, perhaps in terms of developing the capacity necessary to process and analyse data. It could also suggest such databases are not useful for meeting local needs. Bowles-Newark et al. (2015) found a lot of uncertainty among national CBD focal points on the accessibility and applicability of global data sets. Issues of access and perceived relevance may be compounded by the fact that data users need at least 40 types of data (not only species presence or population trends), and that global databases do not exist for all of these variables.

With multiple challenges to data use uncovered by our user survey, multiple options need to be considered to help remove those challenges and make data more readily available. While more funding for monitoring is obviously key and was the main blockage noted by respondents to our survey, other solutions include allocating more of existing budgets to monitoring (Stephenson 2019; Badalotti et al. 2022). Furthermore, some studies suggest that starting biodiversity monitoring programmes in Africa could require as little as US\$ 30,000–50,000 per country per year (Pereira et al. 2010; Wotton et al. 2020). Nonetheless, this will not avoid the fact that, in some cases, other conservation work is deemed more urgent than monitoring. For example, in Kenya there is some evidence that, at least in protected areas, managers often have to prioritize park security, anti-poaching, and the monitoring of illegal activity over the monitoring of species status (Stephenson et al. 2021). This further underlines the resource challenges and decisions facing many managers. The pervasive poverty in East Africa, both in terms of financial income and access to education and health, is a major constraint to the development of conservation and monitoring projects in this region (Kinzig and McShane 2015).

Globally, per capita GDP has been shown to be correlated with the number of species monitoring programmes in each country (Moussy et al. 2022) but GDP did not have a significant relationship with data availability in East Africa. However, our study found a positive correlation between per capita international tourism income and the number of priority species and species with data in GBIF and the LPI. Africa is a leading destination for nature and wildlife experiences (Higginbottom 2004) and the most established tourism products in Africa are wildlife related, such as safaris for the “Big Five” mammals, gorilla tourism, birdwatching and scuba diving (World Tourism Organization 2014). As discussed by Nyaupane and Poudel (2011), tourism development generates income for conservation and conversely biodiversity conservation makes places attractive to tourists. Although we can only infer causality, tourism revenue may explain better than GDP the number of priority species and the availability of data in East Africa because of the wildlife focus of so much tourism and the investments made in conserving species that attract those tourists. It would be interesting to conduct more detailed case studies and explore the links between tourism and data availability within and across regions.

Our findings underline the point raised by others that, given the high levels of species endemism and diversity and the low levels of GDP in sub-Saharan Africa, it is especially important for the international conservation community of donors and NGOs to ramp up support for species monitoring (Stephenson et al. 2017a, 2020, 2021). The second most important challenge for data users in East Africa was the use of technology, yet the use of remote sensing and other modern techniques such as DNA-based approaches will be essential for improving species monitoring across Africa (Stephenson et al. 2020, 2021). More guidance and capacity building support for monitoring should therefore be provided to project managers (Stephenson et al. 2015b, 2017a, b; Schmeller et al. 2017; Stephenson 2019; Badalotti et al. 2022). Capacity building would be further enhanced if conservationists implemented more pilots and case studies on using global data for national reporting

that are then communicated to help share lessons (Bowles-Newark et al. 2015; Stephenson et al. 2015b).

Regional schemes for biodiversity monitoring that should be promoted and supported in order to fill data gaps include the Global Coral Reef Monitoring Network (Obura et al. 2017), and the GlobWetland Africa Project (Gardner et al. 2015). As well as sharing data with global databases, regional databases in Africa should also be expanded, including FishBase for Africa (<http://www.fishbaseforafrica.org/>), the Albertine Rift Conservation Society Biodiversity Management Information System (<http://arbmis.arcosnetwork.org/>), and the Africa Marine Atlas and African portal on the Ocean Biogeographic Information System (<http://www.iobis.org/>). The GBIF Secretariat (2019) is helping African countries to create networks of data holders and users and to digitize existing data. National efforts should also be ramped up to collect, use and share data, building on successful models like SANBI (2022) and the Endangered Wildlife Trust (2022).

Conclusions

East African biodiversity is under threat and in need of successful conservation action to preserve species, habitats and the ecosystem services for current and future generations. However, effective conservation action and sustainable development requires data for adaptive management and, as we demonstrated, this poses severe challenges. While global databases provide data for most priority species, taxonomic and geographic biases exist. Furthermore, many conservationists face capacity and resource challenges in accessing and using the data they need—and the data decision makers need most are the data that are hardest for them to access.

Based on our findings, we propose a series of actions to enhance data availability for key decision makers. Priorities include: the development by governments and their academic and NGO partners of long-term monitoring programmes for priority species (taking into account the need to counter identified data biases); the mobilisation of more financing and a larger proportion of existing conservation financing for species monitoring; the development of capacity for data collection, including for the use of the latest remote sensing and DNA technologies; increased engagement of citizen scientists to help governments and NGOs collect data; improved data sharing between national and global databases; and improved communications and case studies on the accessibility and uses of global databases for national monitoring.

These actions will require collaboration between governments and civil society, and more support from wealthier countries to those with more biodiversity. If this support is forthcoming, and the motivation to demonstrate national contributions towards the post-2020 global biodiversity framework and the Sustainable Development Goals helps create the appropriate enabling environment, we hope to see an increase in the capacity of East African states to collect, share and use biodiversity data and enhance the impact of conservation action on the ground.

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1007/s10531-022-02497-4>.

Acknowledgements We would like to thank the following organisations for participating in the study: African Parks, Akagera Management Company, Albertine Rift Conservation Society, Association Burundaise pour la protection de la Nature, The Aspinall Foundation, Big Life Foundation, Center of Excellence in Biodiversity and Natural Resource Management, University of Rwanda, Chimpanzee Trust, Dahari,

Endangered Wildlife Trust, Environment Affairs Department Malawi, Ethiopian Wildlife and Natural History Society, Fondation Segre, Giraffe Conservation Foundation, International Union for the Conservation of Nature, Lemur Conservation Association, Lemur Conservation Foundation, Lemur Love, Madagasikara Voakajy, Ministère de l'Agriculture, de la Pêche et de l'Environnement, Union des Comores, Nature Kenya, Nature Rwanda, Nature Uganda, Pamoja Environmental Focus, Sea Sense, Somali Wildlife & Natural History Society, TRAFFIC, Wildlife and Environmental Society of Malawi, Wildlife Action Group, Wildlife Works, Zoological Society of London. We also thank Estelle Milliet for providing advice on the questionnaire survey, Julie Guenat for useful feedback on an earlier draft, and the feedback from two anonymous referees. PJS was part-funded by the Swiss Network for International Studies (<https://snis.ch>).

Author contributions PJS and IM conceptualised the study and designed the methodology; IM collected the data; IM and JG analysed and visualized the data; IM and PJS drafted the manuscript; all authors reviewed and edited the final manuscript; LF and PJS were responsible for project administration and supervision.

Funding Open access funding provided by the University of Lausanne. This study was part funded by a grant from the Swiss Network for International Studies.

Data availability Where data user agreements of the data sources allow, data have been made available as Supplementary Material in the annex.

Declarations

Conflict of interest The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Ethical approval All work conforms to the legal requirements of the country in which it was carried out, including those relating to conservation and welfare, and to the Journal's policy on these matters. All survey participants were informed of the aims of the work and that their answers would be used in data analysis.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Albert C, Luque GM, Courchamp F (2018) The twenty most charismatic species. *PLoS ONE* 13(7):1–12
- Amano T, Sutherland W (2013) Four barriers to the global understanding of biodiversity conservation: wealth, language, geographical location and security. *Proc R Soc B: Biol Sci* 280(1756):1–7
- Amano T, Lamming JDL, Sutherland WJ (2016) Spatial gaps in global biodiversity information and the role of citizen science. *Bioscience* 66(5):393–400
- Bachman SP, Nic Lughadha EM, Rivers MC (2018) Quantifying progress toward a conservation assessment for all plants. *Conserv Biol* 32(3):516–524
- Badalotti A, van Galen L, Vié J-C, Stephenson PJ (2022) Improving the monitoring of conservation programmes: lessons from a grant-making initiative for threatened species. *Oryx* 56(2):288–294
- Balmford A (2003) Conservation planning in the real world: South Africa shows the way. *Trends Ecol Evol* 18:435–438
- Bhatt R, Gill MJ, Hamilton H, Han X, Linden HM, Young BE (2020) Uneven use of biodiversity indicators in 5th national reports to the convention on biological diversity. *Environ Conserv* 47(1):15–21
- Bonnet X, Shine R, Lourdaux O (2002) Taxonomic chauvinism. *Trends Ecol Evol* 17(1):1–3
- Bowles-Newark NJ, Chenery A, Misrachi M, Despot-Belmonte K (2015) Barriers to the Use of global indicators and datasets to support NBSAP implementation and national reporting processes. UNEP-WCMC, Cambridge

- Brooks TM, Butchart SH, Cox NA, Heath M, Hilton-Taylor C, Hoffmann M, Kingston N, Rodríguez JP, Stuart SN, Smart J (2015) Harnessing biodiversity and conservation knowledge products to track the Aichi Targets and Sustainable Development Goals. *Biodiversity* 16(2–3):157–174
- Brooks M, Kristensen K, van Benthem K, Magnusson A, Berg CW, Nielsen A, Skaug HJ, Mächler M, Bolker B (2017) GlmmTMB balances speed and flexibility among packages for zero-inflated generalized linear mixed modeling. *R J* 9:378–400
- Bubb P, Chenery A, Herkenrath P, Kapos V, Mapendembe A, Stanwell-Smith D, Walpole M (2011) National indicators, monitoring and reporting for the strategic plan for biodiversity 2011–2020: a review of experience and recommendations in support of the CBD ad hoc technical expert group (AHTEG) on indicators for the strategic plan 2011–2020. UNEP-WCMC, Cambridge
- Cardoso P, Leather SR (2019) Predicting a global insect apocalypse. *Insect Conserv Divers* 12:263–267
- Cardoso P, Erwin TL, Borges PA, New TR (2011) The seven impediments in invertebrate conservation and how to overcome them. *Biol Conserv* 144:2647–2655
- Chandler M, See L, Copas K, Bonde AM, López BC, Danielsen F, Legind JK, Masinde S, Miller-Rushing AJ, Newman G, Rosemartin A, Turak E (2017) Contribution of citizen science towards international biodiversity monitoring. *Biol Conserv* 213:280–294
- Chen C, Quan R, Cao G, Yang H, Burton AC, Meitner M, Brodie JF (2019) Effects of law enforcement and community outreach on mammal diversity in a biodiversity hotspot. *Conserv Biol* 33(3):612–622. <https://doi.org/10.1111/cobi.13232>
- Christie AP, Amamo T, Martin PA, Petrovan SO, Shackelford GE, Simmons BI, Smith RK, Williams DR, Wordley CFR, Sutherland WJ (2021) The challenge of biased evidence in conservation. *Conserv Biol* 35(1):249–262
- Collen B, Loh J, Whitmee S, McRae L, Amin R, Baillie JEM (2009) Monitoring change in vertebrate abundance: the Living Planet Index. *Conserv Biol* 23(2):317–327. <https://doi.org/10.1111/j.1523-1739.2008.01117.x>
- Colléony A, Clayton S, Couvet D, Saint Jalme M, Prévot A-C (2017) Human preferences for species conservation: animal charisma trumps endangered status. *Biol Conserv* 206:263–269
- Convention on Biological Diversity (2010) The strategic plan for biodiversity 2011–2020 and the Aichi Biodiversity Targets. UNEP/CBD/COP/DEC/X/2
- Convention on Biological Diversity (2021) National Reports. Website <https://www.cbd.int/reports/>. Accessed 30 May 2021
- Costello MJ (2009) Motivating online publication of data. *Bioscience* 59(5):418–427
- Costello MJ, Michener WK, Gahegan M, Zhang ZQ, Bourne PE (2013) Biodiversity data should be published, cited, and peer reviewed. *Trends Ecol Evol* 28(8):454–461
- Craigie ID, Baillie JE, Balmford A, Carbone C, Collen B, Green RE, Hutton JM (2010) Large mammal population declines in Africa's protected areas. *Biol Conserv* 143:2221–2228
- Danovaro R, Fanelli E, Aguzzi J, Billett D, Carugati L, Corinaldesi C, Dell'Anno A, Gjerde K, Jamieson AJ, Kark S, McClain C, Levin L, Levin N, Ramirez-Llodra E, Ruhl H, Smith CR, Snelgrove PVR, Thomsen L, Van Dover CL, Yasuhara M (2020) Ecological variables for developing a global deep-ocean monitoring and conservation strategy. *Nat Ecol Evol* 4:181–192. <https://doi.org/10.1038/s41559-019-1091-z>
- dos Santos JW, Correia RA, Malhado ACM, Campos-Silva JV, Teles D, Jepson P, Ladle RJ (2020) Drivers of taxonomic bias in conservation research: a global analysis of terrestrial mammals. *Anim Conserv* 23(6):679–688
- Endangered Wildlife Trust (2022) Biodiversity Data. <https://www.ewt.org.za/resources/resources-biodiversity-data/>. Accessed 10 Mar 2022
- FAO (2021) FAOLEX Database. <https://www.fao.org/faolex/en/>. Accessed 12 May 2021
- Fitzpatrick U, Murray ETE, Paxton RJ, Brown MJF (2007) Building on IUCN regional red lists to produce lists of species of conservation priority: a model with Irish bees. *Conserv Biol* 21:1324–1332
- Fox J, Weisberg S (2019) *An R companion to applied regression*, 3rd edn. Sage, Thousand Oaks, CA
- Gardner RC, Barchiesi S, Beltrame C, Finlayson CM, Galewski T, Harrison I, Paganini M, Perennou C, Pritchard D, Rosenqvist A, Walpole M (2015) State of the World's Wetlands and their Services to People: A compilation of Recent Analyses. Ramsar Briefing Note No. 7. Ramsar Convention Secretariat, Gland
- Garibaldi L (2014) The value of African fisheries. Fishery and aquaculture economics and policy division. FAO, Rome
- Gascon C, Brooks TM, Contreras-MacBeath T, Heard N, Konstant W, Lamoreux J, Launay F, Maunder M, Mittermeier RA, Molur S, Al Mubarak RK (2015) The importance and benefits of species. *Curr Biol* 25(10):R431–R438. <https://doi.org/10.1016/j.cub.2015.03.041>
- GBIF.org (2021). GBIF Home Page. <https://www.gbif.org>. Accessed 1 Aug 2021

- Secretariat GBIF (2019) Establishing an effective GBIF participant node: concepts and general considerations. GBIF Secretariat, Copenhagen. <https://doi.org/10.15468/doc-z79c-sa53>
- Gonçalves SC, Haelewaters D, Furci G, Meller GM (2021) Include all fungi in biodiversity goals. *Science* 373(6553):403–403
- Han X, Smyth RL, Young BE, Brooks TM, de Lozada AS, Bubbs P, Butchart SHM, Larsen FW, Hamilton H, Hansen MC, Turner WR (2014) A biodiversity indicators dashboard: addressing challenges to monitoring progress towards the Aichi Biodiversity Targets using disaggregated global data. *PLoS ONE* 9:e112046
- Hartig F (2021) DHARMA: residual diagnostics for hierarchical (multi-Level/mixed) Regression Models. <https://CRAN.R-project.org/package=DHARMA>. Accessed 30 Oct 2021
- Higginbottom K (2004) *Wildlife tourism: impacts, management and planning*. Common Ground Publishing, Altona, VIC
- Hochkirch A, Samways MJ, Gerlach J, Böhm M, Williams P, Cardoso P, Cumberlidge N, Stephenson PJ, Seddon MB, Clausnitzer V, Borges PAV, Mueller GM, Pearce-Kelly P, Raimondo DC, Dijkstra K-DB (2021) A strategy for the next decade to address data deficiency in neglected biodiversity. *Conserv Biol* 35(2):502–509
- Hu Y, Luo Z, Chapman CA, Pimm SL, Turvey ST, Lawes MJ, Peres CA, Lee TM, Fan P (2019) Regional scientific research benefits threatened-species conservation. *Natl Sci Rev* 6(6):1076–1079
- Huxley J (1961) *The conservation of wild life and natural habitats in Central and East Africa*. UNESCO, Paris
- International Monetary Fund (2021) IMF data on GDP. <https://www.imf.org/external/datamapper/NGDPD@WEO/OEMDC/ADVEC/WEOWORLD>. Accessed 1 Oct 2021
- IPBES (2018) The IPBES regional assessment report on biodiversity and ecosystem services for Africa. In: Archer E, Dziba L, Mulongoy KJ, Maoela MA, Walters M (eds) Secretariat of the intergovernmental science-policy platform on biodiversity and ecosystem services. IPBES, Bonn
- IUCN (2012) IUCN Red List Categories and Criteria: Version 3.1. IUCN, Gland
- IUCN (2021) The IUCN Red List of Threatened Species. Version 2021–2. <https://www.iucnredlist.org>. Accessed 15 Apr 2021
- IUCN, UNEP, FAO (2021). *Ecolex*. <https://www.ecolex.org/>. Accessed 10 May 2021
- Kinzig AP, McShane TO (2015) Conservation in Africa: exploring the impact of social, economic and political drivers on conservation outcomes. *Environ Res Lett* 10(9):090201. <https://doi.org/10.1088/1748-9326/10/9/090201>
- Krause M, Robinson K (2017) Charismatic species and beyond: how cultural schemas and organisational routines shape conservation. *Conserv Soc* 15:313–321
- Le Berre M, Noble V, Pires M, Médail F, Diadema K (2019) How to hierarchise species to determine priorities for conservation action? A critical analysis. *Conserv Soc* 28:3051–3071
- Leather SR (2009) Taxonomic chauvinism threatens the future of entomology. *Biologist* 56:10–13
- Living Planet Index (2021) Living Planet Index. <https://www.livingplanetindex.org/home/index>. Accessed 1 Aug 2021
- Loh J, Green RE, Ricketts T, Lamoreux J, Jenkins M, Kapos V, Randers J (2005) The Living Planet Index: using species population time series to track trends in biodiversity. *Philos Trans R Soc B: Biol Sci* 360(1454):289–295. <https://doi.org/10.1098/rstb.2004.1584>
- Mace GM, Possingham HP, Leader-Williams N (2007) Prioritizing choices in conservation. In: Macdonald DW, Service K (eds) *Key topics in conservation biology*. Blackwell Publishing, Oxford, pp 17–34
- Mastrángelo ME, Pérez-Harguindeguy N, Enrico L, Bennett E, Lavorel S, Cumming GS, Abeygunawardane D, Amarilla LD, Burkhard B, Egoh BN, Frishkoff L (2019) Key knowledge gaps to achieve global sustainability goals. *Nat Sustain* 2(12):1115–1121
- McRae L, Deinet S, Freeman R (2017) The diversity-weighted living planet index: controlling for taxonomic bias in a global biodiversity indicator. *PLoS ONE* 12(1):1–20
- Meyer MA, Booker JM (2001) *Eliciting and analyzing expert judgment*. Society for Industrial and Applied Mathematics, Philadelphia
- Mittermeier RA, Robles Gil P, Hoffmann M, Pilgrim J, Brooks T, Goettsch Mittermeier C, Lamoreux J, Da Fonseca GAB (2004) Hotspots revisited. Cemex, San Pedro Garza García
- Moussy C, Burfield IJ, Stephenson PJ, Newton AFE, Butchart SHM, Sutherland WJ, Gregory RD, McRae L, Bubbs P, Roesler I, Ursino C, Wu Y, Retief EF, Udin JS, Urazaliyev R, Sanchez-Clavijo LM, Lartey E, Donald PF (2022) A quantitative global review of species population monitoring. *Conserv Biol* 36:e13721
- Nyaupane GP, Poudel S (2011) Linkages among biodiversity, livelihood, and tourism. *Ann Tour Res* 38(4):1344–1366

- Obura DO, Gudka M, Abdou Rabi F, Bacha Gian S, Bijoux J, Freed S, Maharavo J, Mwaura J, Porter S, Sola E, Wickel J (2017) Coral reef status report for the Western Indian Ocean. Global Coral Reef Monitoring Network (GCRMN)/International Coral Reef Initiative (ICRI)
- Olson D, Dinerstein E (2002) The Global 200: priority ecoregions for global conservation. *Ann Mo Bot* 89(2):199–224
- Pereira HM, Belnap J, Brummitt N, Collen B, Ding H, Gonzalez-Espinosa M, Gregory RD, Honrado J, Jongman RHG, Julliard R, McRae L, Proença V, Rodrigues P, Opige M, Rodriguez JP, Schmeller DS, van Swaay C, Vieira C (2010) Global biodiversity monitoring. *Front Ecol Environ* 8:459–460
- Piepho H-P (2004) An algorithm for a letter-based representation of all-pairwise comparisons. *J Comput Gr Stat* 13:456–466
- Pimm S, Raven P, Peterson A, Şekercioğlu ÇH, Ehrlich PR (2006) Human impacts on the rates of recent, present, and future bird extinctions. *Proc Natl Acad Sci USA* 103(29):10941–10946
- R Core Team (2021) R: a language and environment for statistical computing. Version 4.1.1. R Foundation for Statistical Computing, Vienna
- Reichman OJ, Jones MB, Schildhauer MP (2011) Challenges and opportunities of open data in ecology. *Science* 331(6018):703–705
- Robinson NM, Scheele BC, Legge S, Southwell DM, Carter O, Lintermans M, Radford JQ, Skroblin A, Dickman CR, Koleck J, Wayne AF (2018) How to ensure threatened species monitoring leads to threatened species conservation. *Ecol Manage Restor* 19(3):222–229
- Rounsevell MDA, Harfoot M, Harrison PA, Newbold T, Gregory RD, Mace GM (2020) A biodiversity target based on species extinctions. *Science* 368(6496):1193–1195
- SANBI (2022) Infobases. <https://www.sanbi.org/resources/infobases/>. Accessed 31 Mar 2022
- Sanders MJ, Miller L, Bhagwat SA, Rogers A (2021) Conservation conversations: a typology of barriers to conservation success. *Oryx* 55(2):245–254
- Schmeller DS, Böhm M, Arvanitidis C, Barber-Meyer S, Brummitt N, Chandler M, Chatzinikolaou E, Costello MJ, Ding H, García-Moreno J, Gill M (2017) Building capacity in biodiversity monitoring at the global scale. *Conserv Soc* 26(12):2765–2790
- Seto KC, Güneralp B, Hutyra LR (2012) Global forecasts of urban expansion to 2030 and direct impacts on biodiversity and carbon pools. *Proc Natl Acad Sci USA* 109(40):16083–16088
- Sievers M, Brown CJ, Tulloch VJD, Pearson RM, Haig JA, Turschwell MP, Connolly RM (2019) The role of vegetated coastal wetlands for marine megafauna conservation. *Trends Ecol Evol* 34(9):807–817
- Sitati NW, Ipara H (2012) Indigenous ecological knowledge of a human–elephant interaction in Transmara District, Kenya: implications for research and management. *Adv Anthropol* 2(3):107–111
- Stephenson PJ (2019) The Holy Grail of biodiversity conservation management: monitoring impact in projects and project portfolios. *Perspect Ecol Conserv* 17(4):182–192
- Stephenson PJ, Ntiamao-Baidu Y (2010) Conservation planning for a widespread threatened species: WWF and the African elephant *Loxodonta africana*. *Oryx* 44:194–204
- Stephenson PJ, Stengel C (2020) An inventory of biodiversity data sources for conservation monitoring. *PLoS ONE* 15:e0242923
- Stephenson PJ, Burgess ND, Jungmann L, Loh J, O'Connor S, Oldfield T, Reidhead W, Shapiro A (2015a) Overcoming the challenges to conservation monitoring: integrating data from in situ reporting and global data sets to measure impact and performance. *Biodiversity* 16:68–85
- Stephenson PJ, O'Connor S, Reidhead W, Loh J (2015b) Using biodiversity indicators for conservation. *Oryx* 49:396
- Stephenson PJ, Bowles-Newark N, Regan E, Stanwell-Smith D, Diagana M, Hoft R, Abarchi H, Abrahamse T, Akello C, Allison H, Banki O, Batiemo B, Dieme S, Domingos A, Galt R, Githaiga CW, Guindol AB, Hafashimana DLN, Hirsch T, Hobern D, Kaaya J, Kaggwa R, Kalemba MM, Linjouom I, Manaka B, Mbwambo Z, Musasa M, Okoree E, Rwetsiba A, Siams AB, Thiombiano A (2017a) Unblocking the flow of biodiversity data for decision-making in Africa. *Biol Conserv* 213:335–340
- Stephenson PJ, Brooks TM, Butchart SHM, Fegraus E, Geller GN, Hoft R, Hutton J, Kingston N, Long B, McRae L (2017b) Priorities for big biodiversity data. *Front Ecol Environ* 15:124–125
- Stephenson PJ, Ntiamao-Baidu Y, Simaika JP (2020) The use of traditional and modern tools for monitoring wetlands biodiversity in Africa: challenges and opportunities. *Front Environ Sci* 8:61
- Stephenson PJ, Bakarr M, Bowles-Newark N, Kleinschroth F, Mapendembe A, Ntiamao-Baidu Y, Obura D, Ratsifandrihamana N, Simaika J, Sitati N, Sumbi P, Teferi T (2021) Conservation science in Africa: mainstreaming biodiversity information into decision-making. In: Ferreira CC, Klütsch CFC (eds) *Closing the Knowledge-Implementation Gap in Conservation Science—evidence transfer across spatiotemporal scales and different stakeholders*. Wildlife Research Monograph Number 4. Springer, New York, pp 287–321. https://doi.org/10.1007/978-3-030-81085-6_11

- Stephenson PJ, Londoño-Murcia MC, Borges PAV, Claassens L, Frisch-Nwakanma H, Ling N, McMullan-Fisher S, Meeuwig JJ, Unter KMM, Walls JL, Burfield IJ, Correa DCV, Geller GN, Montenegro Paredes I, Mubalama LK, Ntiamao-Baidu Y, Roesler I, Rovero F, Sharma YP, Wisesa Wiwardhana N, Yang J, Fumagalli L (2022) Measuring the impact of conservation: the growing importance of monitoring fauna, flora and funga. *Diversity* 14:824
- Tenopir C, Allard S, Douglass K, Aydinoglu AU, Wu L, Read E, Manoff M, Frame M (2011) Data sharing by scientists: practices and perceptions. *PLoS ONE* 6:e21101
- Thapa I, Butchart SHM, Gurung H, Stattersfield AJ, Thomas DHL, Birch JC (2014) Using information on ecosystem services in Nepal to inform biodiversity conservation and local to national decision-making. *Oryx* 50:147–155
- Theobald EJ, Ettinger AK, Burgess HK, DeBey LB, Schmidt NR, Froehlich HE, Wagner C, HilleRisLambers J, Tewksbury J, Harsch MA, Parrish JK (2015) Global change and local solutions: tapping the unrealized potential of citizen science for biodiversity research. *Biol Conserv* 181:236–244
- Titley MA, Snaddon JL, Turner EC (2017) Scientific research on animal biodiversity is systematically biased towards vertebrates and temperate regions. *PLoS ONE* 12:e0189577
- Tittensor DP, Walpole M, Hill SL, Boyce DG, Britten GL, Burgess ND, Butchart SHM, Leadley PW, Regan EC, Alkemade R, Baumung R, Bellard C, Bouwman L, Bowles-Newark NJ, Chenery AM, Cheung WWL, Christensen V, Cooper HD, Crowther AR, Dixon MJR, Galli A, Gaveau V, Gregory RD, Gutierrez NL, Hirsch TL, Höft R, Januchowski-Hartley SR, Karmann M, Krug CB, Leverington FJ, Loh J, Lojenga RK, Malsch K, Marques A, Morgan DHW, Mumby PJ, Newbold T, Noonan-Mooney K, Pagad SN, Parks BC, Pereira HM, Robertson T, Rondinini C, Santini L, Scharlemann JPW, Schindler S, Sumaila UR, Teh LSL, van Kolck J, Visconti P, Ye Y (2014) A mid-term analysis of progress toward international biodiversity targets. *Science* 346:241–244
- Troudet J, Grandcolas P, Blin A, Vignes-Lebbe R, Legendre F (2017) Taxonomic bias in biodiversity data and societal preferences. *Sci Rep* 7(1):1–14
- UNEP-WCMC (2021) Species+. <https://www.speciesplus.net/>. Accessed 15 May 2021
- UNWTO (2021) World Tourism Organization eLibrary. <https://www.e-unwto.org>. Accessed 1 Oct 2021
- van Strien AJ, Gmelig Meyling AW, Herder JE, Hollander H, Kalkman VJ, Poot MJM, Turnhout S, van der Hoorn B, van Strien-van LWFH, van Swaay CAM, van Turnhout CAM, Verweij RJT, Oerlemans NJ (2016) Modest recovery of biodiversity in a western European country: The Living Planet Index for the Netherlands. *Biol Conserv* 200:44–50
- VanRossum G, Drake FL (2009) Python 3 reference manual. CreateSpace, Scotts Valley, CA
- Wilson HB, Joseph LN, Moore AL, Possingham HP (2011) When should we save the most endangered species? *Ecol Lett* 14:886–890
- World Tourism Organization (2014) Towards measuring the economic value of wildlife watching tourism in Africa – Briefing Paper. UNWTO, Madrid
- Wotton SR, Eaton MA, Sheehan D, Munyekenye FB, Burfield IJ, Butchart SHM, Moleofi K, Nalwanga-Wabwire D, Pomeroy D, Senyato KJ, Gregory RD (2020) Developing biodiversity indicators for African birds. *Oryx* 54:62–73
- WWF (2020) Living planet report 2020: bending the curve of biodiversity loss. In: Almond REA, Grooten M, Petersen T (eds) Living planet report 2020: bending the curve of biodiversity loss. WWF, Gland
- WWF-Canada (2020) Living planet report Canada: wildlife at risk. In: Currie J, Snider J, Giles E (eds) Living planet report Canada: wildlife at risk. World Wildlife Fund Canada, Toronto

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Authors and Affiliations

Inès Moreno^{1,2,3} · Jérôme Marcel Walter Gippet⁴  · Luca Fumagalli¹  · P. J. Stephenson^{1,2} 

Inès Moreno
ines.moreno@bluewin.ch

Jérôme Marcel Walter Gippet
Jerome.Gippet@unil.ch

Luca Fumagalli
Luca.Fumagalli@unil.ch

- ¹ Laboratory for Conservation Biology, Department of Ecology & Evolution, University of Lausanne, CH-1015 Lausanne, Switzerland
- ² IUCN SSC Species Monitoring Specialist Group, Lausanne, Switzerland
- ³ Present Address: KORA, CH-3063 Ittigen, Switzerland
- ⁴ Department of Ecology & Evolution, University of Lausanne, CH-1015 Lausanne, Switzerland